



To evaluate the influence of inhaled bronchodilator therapy on spirometric test variables in obese and non-obese asthmatics

Dr. Latha Sarma^{1*}, Nandan Putti^{1*}, Dr. Avinash Chaskar¹, Dr. Kapil Alias Mohit Chilana¹ and Asmath Qureshi¹

¹ Department of Pulmonology, Krishna Institute of Medical Sciences, Minister Road, Secunderabad-500003, Telangana, India.

Abstract

In India, the prevalence of overweight individuals is reported to be between 20-40% in various studies, with over 5-10% being obese. Obesity acts as a risk factor for asthma through various mechanisms and it also reduces the spirometric variables as well as total lung capacity and functional residual capacity. Forced expiratory volume in one second (FEV₁) was done on 100 patients in tertiary care hospital using RMS Helios 401 electronic spirometer, by Recorders and Medicare Systems Pvt. Ltd. All the statistical analysis was done using SPSS, Ver. 17.0, (IBM, Chicago, Illinois and student unpaired T-test and analysis of variance (ANOVA) test was used for estimating results. 59 were males and 41 were females, of the 59 males, 29 were obese and 30 were non-obese and of the 41 females 20 were obese and 21 were non-obese. There was highly significant difference found in the spirometric variables in obese and non-obese asthmatics. There is significant amount of reversibility (% change) after inhalation of short acting beta-2 agonist in non-obese group as compared with obese group with mean percentage change in non-obese group being 16.20% and obese group 13.58%. p value 0.002 (p<0.05). The amount of reversibility (mean % change) was found to be more for controlled asthmatic groups (16.88%) as compared with partly controlled asthmatics (14.84%) and uncontrolled asthmatic groups (12.78%) p value 0.001 (p<0.05). To conclude, it can be said that age does not have any significant influence on the levels of asthma control in both obese and non-obese asthmatics. Obesity has a significant influence on various spirometric variables.

Keywords: asthma; obese; non-obese; spirometry values; reversibility; cross sectional study

***Corresponding authors:** Dr. Latha Sarma, Critical respiratory & Sleep specialist, Department of Pulmonology, Krishna Institute of Medical Sciences, Minister Road, Secunderabad-500003, Telangana, India, Email: dr.lathasarma@gmail.com and Nandan Putti, Department of Pulmonology, Krishna Institute of Medical Sciences, Minister Road, Secunderabad-500003, Telangana, India. Tel.: +919014036776; Email: naddyn7@gmail.com

Received 20 June 2015; Revised 3 September 2015; Accepted 10 September 2015

Citation: Latha Sarma, Nandan Putti, Avinash Chaskar, Mohit Chilana KA, Asmath Qureshi. To evaluate the influence of inhaled bronchodilator therapy on spirometric test variables in obese and non-obese asthmatics. J Med Sci Res. 2015; 3(4):166-171. DOI: <http://dx.doi.org/10.17727/JMSR.2015/3-032>

Copyright: © 2015 Latha Sarma, et al. Published by KIMS Foundation and Research Center. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

In the present era of health consciousness and better educational standards, health, lifestyle and behaviour of an individual play a key role. Asthma is a chronic disorder affecting millions of people worldwide. The prevalence of asthma is around 300 million and is expected to increase another 100 million by 2025 (WHO: Global surveillance, prevention and control of chronic respiratory diseases; 2007) [1]. In India, the prevalence of overweight individuals is reported to be between 20-40% in various studies, with over 5-10% being obese [2, 3].

There is epidemiological data indicating a causal relationship between obesity and asthma. Several

studies suggest that weight loss and weight gain may have an effect on the clinical course of asthma [2, 4-9]. Prospective studies show that obesity is a risk factor for asthma, though the mechanisms linking the two are not fully understood. Obesity is a disorder associated with systemic inflammation, and an increase in inflammatory mediators, which may also result in airway hyper reactivity, which is observed in obese individuals [4].

Adipose tissue is an important source of cytokines and contributes to the inflammatory milieu. Apart from general obesity, visceral adipose tissue is the key factor in the formation of low grade chronic inflammation in obese individuals [10] and plays a role in remodelling, which is characteristic of asthma. High levels of interleukin-6 (IL-6) and tumour necrosis factor-alpha (TNF- α) are observed in individuals with visceral obesity. IL-6 is the primary mediator in acute phase response and leads to the synthesis of C-reactive protein(CRP), which exacerbates the inflammatory response [11].

Asthma is a chronic inflammatory disease of the airways [12]. Obesity acts as a risk factor for asthma through various mechanisms. TNF- α is expressed in the airways and TNF- α may increase contractility against airway constrictor agents; in other words they may increase airway hyper reactivity [2].

Asthma significantly lowers specific conductance and the spirometric variables, while increasing airway reactivity and residual volume. Obesity also reduces the spirometric variables as well as total lung capacity and functional residual capacity. Residual volume, specific conductance, and airway responsivity are unaltered [13]. Besides, it may reduce respiratory muscle strength, decrease thoracic cage compliance and impede diaphragmatic excursion, especially when it is massive and central. Thus, there is over diagnosing asthma in the obese. Asthma has been reported to be more difficult to control in obese patients compared to individuals with normal weight [14-17].

Therefore, in the light of this new information, further studies are required to explore the relation between asthma and obesity in India.

Materials and methods

A cross-sectional study was conducted at tertiary care centre Krishna Institute of Medical Sciences,

Ltd. Secunderabad. Over 6 months period on 100 bronchial asthma patients were distributed into two groups, 50 obese and 50 non-obese. Patients who were diagnosed with asthma and fall in the age group of 18-65 years were included in this study.

To limit the study to the effects of asthma and obesity, subjects with history of lung disease other than asthma, like coronary artery disease, congestive heart failure or cor pulmonale and smoking were excluded.

Selected patients were explained the purpose of the study and the need of co-operation were emphasized. All subjects participated in the study voluntarily. Written informed consent was obtained from all patients. Ethical and scientific clearance for the study was obtained from the institutional Ethical Committee.

Spirometry was done on all the subjects using RMS Helios 401 electronic spirometer (Figure 1), by Recorders and Medicare Systems Pvt. Ltd., which is a precalibrated and computerized spirometer. After instruction, each subject performed a minimum of three forced expiratory manoeuvres and following spirometric parameter was recorded for analysis: 1) forced vital capacity (FVC); 2) forced expiratory volume in one second (FEV₁); 3) FEV₁/FVC.



Figure 1: Showing the technique of performing pulmonary function test (PFT) using Helios spirometer.

Body mass index (BMI): Body mass index is an index of weight for height, which is commonly used in classifying overweight and obesity in adult population (Table 1).

BMI= weight in kg/square of height in meters [Quetelet's index].

Table 1: BMI classification.

<i>Classification</i>	<i>BMI</i>
Underweight	< 18.5
Normal range	18.5-22.9
Overweight at risk	23-24.9
Obesity I	25-29.9
Obesity II	≥ 30.

Table 2: The level of asthma control is evaluated by monitoring symptoms and PFT: (Global initiative for asthma (GINA) updated 2012).

<i>Characteristic</i>	<i>Controlled (all of the following)</i>	<i>Partially controlled (any measure present)</i>	<i>Uncontrolled</i>
Daytime symptoms	None (twice/ <per week)	More than twice per week	
Limitation of activities (Including exercise)	None	Any	Three or more features of partly controlled asthma present in any week.
Nocturnal awakenings/ symptoms	None	Any	
Need for rescue treatment	None (twice/ <per week)	More than twice/ week	
Lung function (PEF or FEV ₁)	Normal	<80% predicted or personal best if known	

through spacer. The degree of reversibility in FEV₁ which indicates a diagnosis of asthma which is 12% or 200 ml from pre-bronchodilator value was taken into consideration.

Statistics: All the statistical analysis was done using SPSS, Ver. 17.0, (IBM, Chicago, Illinois). Master sheet of data was made with microsoft excel, graphs and charts were made with MS excel and word. Student unpaired 'T' test was used for comparing the spirometric variables in obese and non-obese asthmatic groups. Each individual quantitative data such as BMI, weight, and duration of asthma are compared considering the levels of asthma using analysis of variance (ANOVA) test.

Results

Out of 100 asthmatics that were enrolled in the study, 59 were males and 41 were females. Of the 59 males, 29 were obese and 30 were non-obese. Of the 41 females 20 were obese and 21 were non-obese. The mean age of obese group is 45.92 years; and non-obese group is 40.78 years. Details are given in figure 2.

The number of patients with uncontrolled asthma was 27 and the mean age was around 43 in the three groups of asthma control as seen in table 3.

Spirometry is the recommended method to establish a diagnosis of asthma, while PEF measurement is more useful for monitoring of asthma (Table 2). Spirometry is reproducible but effort dependent.

Reversibility: The term is applied to rapid improvement in FEV₁ or PEF, measured within minutes after inhalation of a rapid acting β₂ bronchodilator, salbutamol 200 mcg inhaler 2 puffs

Table 3: Age and levels of asthma control.

<i>Levels of asthma control</i>	<i>N</i>	<i>Mean age (Years)</i>
Controlled	30	41.6
Partly controlled	43	43.83
Uncontrolled	27	44.51

As p value is 0.000 (p <0.001) this shows BMI increased from 22.34 to 29.84 the level of asthma control deteriorated from controlled state to uncontrolled state. Detailed data is given in table 4.

Table 4: BMI and levels of asthma control.

<i>Level of asthma control</i>	<i>N</i>	<i>Mean BMI (Kg/m²)</i>
Controlled	30	22.34
Partly controlled	43	24.27
Uncontrolled	27	29.85

The mean percentage change in reversibility of spirometric values in non-obese population was 16.2% whereas in obese population it was 13.58% (Figure 3).





Figure 2: The age and sex distribution of males and females in obese and non-obese group.

Table 5: Comparing the spirometric test results in obese and non-obese asthmatics.

	Class	N	Mean	Std. Deviation on	t value	p value
Pre FEV ₁	Non-obese	50	2.73	0.674	5.203	0.000*
	Obese	50	2.07	0.588		
Pre FVC	Non-obese	50	4.08	0.863	4.126	0.000*
	Obese	50	3.44	0.698		
Pre FEV ₁ /FVC	Non-obese	50	65.96	3.833	5.485	0.000*
	Obese	50	58.64	8.623		
Post FEV ₁	Non-obese	50	3.17	0.818	5.423	0.000*
	Obese	50	2.35	0.687		
Post FVC	Non-obese	50	3.90	0.845	5.047	0.000*
	Obese	50	3.09	0.753		
Post FEV ₁ /FVC	Non-obese	50	80.80	6.302	3.663	0.000*
	Obese	50	75.42	8.256		

Abbreviations: * = indicates high significance.



As the p value is 0.000 ($p < 0.001$) there was highly significant difference found in the spirometric variables in Obese and non-obese asthmatics. The details are given in table 5. The amount of reversibility was the highest in well controlled asthmatics with 16.88% and least in uncontrolled asthmatics with 12.78%. The details are given in table 6.

Discussion

In our study, the comparison of various spirometric variables (FEV_1 , FVC, and FEV_1/FVC) in 100 asthmatics patients randomly divided into 50 obese and 50 non-obese groups, and the influence of inhaled bronchodilator therapy on spirometric test values (i.e. the amount of reversibility) in obese and non-obese asthmatics was analysed.

In our study, the mean age in years for controlled group is 41.6 years, for partly controlled is 43.83 years and for uncontrolled group is 44.51 years, and no statistical significant found between three groups of levels of asthma control considering age, as p value is 0.615 ($p > 0.05$) i.e. variation observed in three groups is by chance and not due to variation in age of subjects. So age does not have any effect on levels of asthma control. So no adjustment of age was done for comparing levels of asthma control. In a study by Lang JE, et al. [18], age is a significant effect modifier on the relationship between obesity and asthma phenotype.

In our study as the mean BMI (kg/m^2) increased from 22.34 to 29.84 the levels of asthma control deteriorated from controlled state to uncontrolled state ($p = 0.000$, highly significant as $p < 0.001$). Studies supporting our study, Mosen D.M. et al. [19] showed

obese adults were more likely than those with normal BMIs ($< 25 kg/m^2$) to report poor asthma-specific quality of life (odds ratio [OR], 2.8; 95% CI, 1.6-4.9), poor asthma control (OR, 2.7; 95% CI, 1.7-4.3), and a history of asthma-related hospitalizations (OR, 4.6; 95% CI, 1.4-14.4). In a case series, by Hernandez LV, et al. [20]. The relation between body mass index and asthma severity is evident; patients with BMI greater than $25 kg/m^2$ are more susceptible to suffer asthma from moderate to severe degree, mainly woman. Obesity seems to be more prevalent among asthmatics with poor control [21].

In our study spirometric variables (FEV_1 , FVC, and FEV_1/FVC) are significantly lower in obese asthmatics as compared with non-obese asthmatics. (p value 0.000, highly significant as $p < 0.001$) indicating obesity having a significant correlation on various spirometric parameters and lung function.

Several studies have different results on the effects of obesity and lung function. There was an effect of BMI on the FEV_1/FVC ratio, indicative of airway obstruction. Particularly, in boys, FEV_1/FVC declined with increasing quintiles of BMI, indicating that obesity promotes airways obstruction. A decline in FEV_1/FVC with BMI in children was also reported by Gold, et al. There is significant amount of reversibility (% change) after inhalation of short acting beta-2 agonist in non-obese group as compared with obese group with mean percentage change in non-obese group being 16.20% and obese group 13.58%. p value 0.002 ($p < 0.05$).

The amount of reversibility (mean % change) was found to be more for controlled groups (16.88%) as compared with partly controlled (14.84%) and uncontrolled groups (12.78%). p value 0.001 ($p < 0.05$). Tantisira, et al. [22] noted an inverse relationship between BMI and bronchodilator response: children with higher BMIs had smaller bronchodilator responses.

Conclusion

Measured spirometric variables are significantly lower in obese asthmatics as compared to non-obese asthmatics. Age does not have any significant influence on the levels of asthma control in both obese and non-obese asthmatics. Levels of asthma control deteriorated from controlled state to uncontrolled state as mean BMI (Kg/m^2) increased. The amount

of reversibility (mean percentage change) after inhalation of short acting β_2 agonist is more for non-obese asthmatic group (16.2%) as compared to obese asthmatic group (13.58%). Obesity has a significant influence on various spirometric variables (FEV_1 , FVC, FEV_1/FVC).

Acknowledgements

Acknowledgements are due to the Departments of Pulmonology, Krishna Institute of Medical Sciences (KIMS), Secunderabad.

Conflict of interest

There was no conflict of interest and all the authors were in tune with the work.

References

1. Chinn S. Obesity and asthma. *Paediatr Respir Rev*. 2006; 7(3):223-228.
2. Sidhu S, Kaur A, Prabhjot. Prevalence of overweight and obesity among urban and rural adult females of Punjab. *Anthropol Anz*. 2005; 63(3):341-345.
3. Sood RK, Gupta AK, Ahluwalia SK, Dhadwal D, Sharma RK. An epidemiological study of obesity in Shimla Town. *Indian J Med Sci*. 1996; 50(10):362-364.
4. Rönmark E, Andersson C, Nyström L, Forsberg B, Järholm B, et al. Obesity increases the risk of incident asthma among adults. *Eur Respir J*. 2005; 25(2):282-288.
5. Jarvis D, Chinn S, Potts J, Burney P; European Community Respiratory Health Survey. Association of body mass index with respiratory symptoms and atopy: results from the European Community Respiratory Health Survey. *Clin Exp Allergy*. 2002; 32(6):831-837.
6. Davis G, Patel JA, Gagne DJ. Pulmonary considerations in obesity and the bariatric surgical patient. *Med Clin North Am*. 2007. 91(3): 433-442.
7. Maniscalco M, Zedda A, Faraone S, Cerbone MR, Cristiano S, et al. Weight loss and asthma control in severely obese asthmatic females. *Respir Med*. 2008; 102(1):102-108.
8. Ford ES. The epidemiology of obesity and asthma. *J Allergy Clin Immunol*. 2005; 115(5):897-909.
9. Schaub B, von Mutius E. Obesity and asthma, what are the links? *Curr Opin Allergy Clin Immunol*. 2005; 5(2):185-193.
10. Park HS, Park JY, Yu R. Relationship of obesity and visceral adiposity with serum concentrations of CRP, TNF-alpha and IL-6. *Diabetes Res Clin Pract*. 2005; 69(1):29-35.
11. Schwarzenberg SJ, Sinaiko AR. Obesity and inflammation in children. *Paediatr Respir Rev*. 2006; 7(4):239-246.
12. Bousquet J, Jeffery PK, Busse WW, Johnson M, Vignola AM. Asthma. From bronchoconstriction to airways inflammation and remodeling. *Am J Respir Crit Care Med*. 2000; 161(5):1720-1745.
13. Nicolacakis K, Skowronski ME, Coreno AJ, West E, Nader NZ, et al. Observations on the physiological interactions between obesity and asthma. *J Appl Physiol*. 2008; 105(5):1533-1541.
14. Lavoie KL, Bacon SL, Labrecque M, Cartier A, Ditto B. Higher BMI is associated with worse asthma control and quality of life but not asthma severity. *Respir Med*. 2006; 100(4): 648-657.
15. Saint-Pierre P, Bourdin A, Chanez P, Daures JP, Godard P. Are overweight asthmatics more difficult to control? *Allergy*. 2006; 61(1):79-84.
16. Dixon AE, Shade DM, Cohen RI, Skloot GS, Holbrook JT, et al. Effect of obesity on clinical presentation and response to treatment in asthma. *J Asthma*. 2006; 43(7):553-558.
17. Boulet LP, Franssen E. Influence of obesity on response to fluticasone with or without salmeterol in moderate asthma. *Respir Med*. 2007; 101(11):2240-2247.
18. Mosen DM, Schatz M, Magid DJ, Camargo CA Jr. The relationship between obesity and asthma severity and control in adults. *J Allergy Clin Immunol*. 2008; 122(3):507-511.
19. Lang JE, Hossain J, Dixon AE, Shade D, Wise RA, et al. Does age impact the obese asthma phenotype? Longitudinal asthma control, airway function, and airflow perception among mild persistent asthmatics. *Chest*. 2011; 140(6):1524-1533.
20. Hernandez LV, Segura NHM. Relation between asthma severity and body mass index. Case series. *Rev Alerg Mex*. 2007; 54(4):107-110.
21. Dixon AE, Shade DM, Cohen RI, Skloot GS, Holbrook JT, et al. Effect of obesity on clinical presentation and response to treatment in asthma. *J Asthma*. 2006; 43(7):553-558.
22. Tantisira KG, Litonjua AA, Weiss ST, Fuhlbrigge AL. Association of body mass with pulmonary function in the Childhood Asthma Management Program (CAMP). *Thorax*. 2003; 58:1036-1041.